

Problem 1*10 points*

For each of the following parts, indicate whether the statement is always true or it can be false by clearly writing “True” or “False.” Full credit will be given for selecting the correct logical value (even with no explanation). You are welcome to briefly explain your reasoning for partial credit (in case your choice is wrong). Diagrams are welcome. Throughout the problem, you may assume that A , B , and C are events with $\mathbb{P}[A] > 0$, $\mathbb{P}[B] > 0$, and $\mathbb{P}[C] > 0$.

(a) If $\mathbb{P}[A|B] = \mathbb{P}[B|A]$, then $\mathbb{P}[A] = \mathbb{P}[B]$.

(b) If $A \cap B = \phi$, $A \cap C = \phi$, and $B \cap C = \phi$, then $\mathbb{P}[A \cup B \cup C] = \mathbb{P}[A] + \mathbb{P}[B] + \mathbb{P}[C]$.

(c) If A and B are independent events, then A^c and B^c are independent events.

(d) It is possible that $\mathbb{P}[B|A] > \mathbb{P}[B]$ (for certain choices of A and B).

(e) If $\mathbb{P}[A] = 0.7$, $\mathbb{P}[B] = 0.4$, and $\mathbb{P}[A \cup B] = 0.8$, then $\mathbb{P}[A^c \cap B^c] = 0.2$.

Problem 2*10 points*

For each of the following parts, indicate whether the statement is always true or it can be false by clearly writing “True” or “False.” Full credit will be given for selecting the correct logical value (even with no explanation). You are welcome to briefly explain your reasoning for partial credit (in case your choice is wrong). Diagrams are welcome. Throughout the problem, you may assume that X is a discrete random variable with PMF $P_X(x)$ and CDF $F_X(x)$.

(a) $\mathbb{P}[X^2 > 0] = \mathbb{P}[X > 0]$.

(b) For a Poisson(λ) random variable, $\mathbb{E}[X^2] = \lambda(1 + \lambda)$.

(c) If the range of X is integer-valued, then the mean $\mathbb{E}[X]$ is integer-valued as well.

(d) For a subset B of the range R_X , we have that $P_{X|B}(x) \geq P_X(x)$ for every $x \in B$.

(e) $\mathbb{E}[X] = \mathbb{E}[X|B] \mathbb{P}[X \in B] + \mathbb{E}[X|B^c] \mathbb{P}[X \in B^c]$

Problem 3 Please complete the following quick calculations.

16 points

(a) Let the events B_1, B_2, B_3 be a partition, with $\mathbb{P}[B_1] = 0.4$, $\mathbb{P}[B_2] = 0.1$, and $\mathbb{P}[B_3] = 0.5$. We know that for event A we have $\mathbb{P}[A|B_1] = 0.2$, $\mathbb{P}[A|B_2] = 0.7$, and $\mathbb{P}[A|B_3] = 0.5$. Calculate $\mathbb{P}[A]$ and $\mathbb{P}[B_3|A]$.

(b) Let $\mathbb{P}[A] = 0.4$, $\mathbb{P}[B] = 0.5$, $\mathbb{P}[C] = 0.3$, $\mathbb{P}[A|B] = 0.3$, $\mathbb{P}[A|C] = 0.2$, $\mathbb{P}[B|C] = 0.5$, and $\mathbb{P}[A \cap B|C] = 0.1$. Are A and B independent? Are A and B conditionally independent given C ?


(c) Let Y be Geometric($1/4$). Calculate $\text{Var}[Y + 4]$ and $\mathbb{E}[3Y + 0.2]$.

(d) Let X be Discrete Uniform($1, 3$). Calculate $\mathbb{E}[X^3]$ as well as the conditional expected value $\mathbb{E}[X|B]$ of X given that $\{X \in B\}$ for $B = \{1, 2\}$.

Problem 4

16 points

You are trying to predict whether a new user of your movie recommendation system will like the latest *Avengers* release, “The Avengers Sequel 23.” Let W be the event that the user has a widescreen TV. Let S be the event that the user likes sequels (in general). Let A be the event that the user likes the new *Avengers* movie.



(a) What is the probability that a new user likes sequels **and** likes the new *Avengers* movie?

(b) What is the probability that a new user likes the new *Avengers* movie?

(c) Given that a user has widescreen, what is the probability they **do not** like the new *Avengers*?

(d) Given that a user likes the new *Avengers*, what is the probability they have a widescreen?

Problem 5*16 points*

Consider the following PMF where β is a constant greater than 0.

$$P_X(x) = \begin{cases} 1/8 & x = 1 \\ 1/2 & x = 2 \\ \beta & x = 5 \\ 0 & \text{otherwise.} \end{cases}$$

(a) Find the value of β that makes $P_X(x)$ a valid PMF. Use this specific value for the rest of the problem. (If you cannot figure this out, just pick any value and proceed.)

(b) Calculate $\text{Var}[X]$.

(c) Calculate $\mathbb{E}\left[\frac{1}{X}\right]$.

(d) Calculate the conditional expected value $\mathbb{E}[X|B]$ given that X falls into $B = \{1, 5\}$.

Problem 7

16 points

You are measuring the number of spikes from a neuron in a one-second window. The resulting random variable X is Poisson(λ).

- (a) After careful study, you have determined that the average number of spikes observed from this neuron in one second is $\mathbb{E}[X] = 2$. What is the probability that you see no spikes at all in a one-second window?
- (b) What is the probability that the number of spikes you see in a one-second window is *less than or equal* to average? (Recall from (a) that the average is 2.)
- (c) Calculate $\mathbb{E}[3X^2 + 2X + 1]$.
- (d) Given that the number of spikes in a one-second window is *less than or equal* to average, what is the conditional expected value of X ?